



FURY MANDEVILLA URC INSTRUCTIONS FROM HIDDEN ACRES NURSERY

1. REMOVE CUTTINGS FROM PLASTIC BAG AND WET WITH CLEAN WATER AND SPREAD OUT IN AREA THAT HAS MIST RUNNING AT LEAST EVERY 10 TO 12 MINUTES.
2. WE SUGGEST THAT CUTTINGS ARE STUCK THE DAY THEY ARRIVE OR NEXT DAY FOR BEST SUCCESS.
3. CUTTINGS SHOULD BE CUT WITH 1" OF STEM BELOW NODE AND 2 LEAVES LEFT ON TOP NODE.
4. WE SUGGEST NO ROOTING HORMONE. IN OUR LAB, WE DISCOVERED THAT THEY DIDN'T LIKE TRADITIONAL ROOTING HORMONES ON MARKET.
5. WE MIST CUTTINGS FOR 5 TO 10 SECONDS EVERY 10 TO 12 MINUTES.
6. CUTTINGS WILL NOT ROOT TILL NIGHT TIME TEMPS ARE ABOVE 65 DEGREES AND ROOT BEST AT 70F NIGHT TIME TEMPS
7. YOU SHOULD SEE ROOTS IN 2 TO 4 WEEKS AND WE TAKE OUT OF MIST IN 5 TO 6 WEEKS. 72 CELL LINERS WILL ROOT OUT IN ABOUT 8 WEEKS NORMALLY
8. DURING THE ROOTING TIME IN MIST, REMOVE ANY LEAVES THAT DROP. LEAVES WILL TURN YELLOW AND DROP, THIS IS NORMAL. STEM IS TAKING BACK NUTRIENTS TO STEM. IT IS IMPORTANT TO KEEP CLEANING LEAVES OUT TO REDUCE DISEASE.
9. WE LIQUID FEED 2 TIME A WEEK WITH 20-20-20 AT A RATE OF 8 OUNCES TO 100 GALLONS OF WATER OR 1 TEASPOON OF 20-20-20 TO 2 GALLONS OF WATER.
10. OUR VARIEGATED LEAF VARIETIES, "HEART THROB" AND "FURY FIRE & ICE" LIKE FERTILIZERS WITH 10 TO 12 MONTH TIME RELEASED COATINGS
11. GREEN LEAFED VARIETIES CAN USE FERTILIZERS WITH 8-9 MONTH AND 10 TO 12 MONTH RELEASE COATINGS
12. OUR FURY SERIES MANDEVILLAS, VARIEGATED LEAF VARIETIES, ARE SENSITIVE TO COPPER FUNGICIDE SPRAYS. FUNGICIDES WE USE ARE HERITAGE, DICONIL, CLEARY'S 3336F, DITHANE, PENTATHALON AND MANY OTHERS. IT IS JUST COPPER THAT THEY DON'T LIKE.
13. WE USE MANY TYPES OF INSECTICIDES AND MITICIDES AND HAVE HAD NO NEGATIVE REACTION OR DAMAGE.

10715 Shankhill Road • Sebring, FL 33875 • 1-800-741-4565 • FAX 863-385-9387
Visit Our Website - hantc.com or contact -wes@hantc.com